

All 2-positive linear maps from $M_3(\mathbb{C})$ to $M_3(\mathbb{C})$
are decomposable*

Wai Shing Tang

* Joint work with Y. Yang and D.H. Leung

National University of Singapore

Quantum Information Theory and Related Topics 2016

Ritsumeikan University, 9 September, 2016

- 1 A Conjecture for 2-positive/2-copositive maps in $B(M_3(\mathbb{C}), M_3(\mathbb{C}))$
 - Origins of the Conjecture
 - The Connections
- 2 A Decomposition Theorem for k -positive maps on Matrix Algebras
 - Block Matrix Approach
 - Some Immediate Consequences
- 3 Questions
 - An Algorithm?
 - An Example?
- 4 References

A Corollary for Generalized Choi Maps in Three Dimensional Matrix Algebra

Let $M_n(\mathbb{C})$ be the C^* -algebra of all $n \times n$ matrices over the complex field.

A Corollary for Generalized Choi Maps in Three Dimensional Matrix Algebra

Let $M_n(\mathbb{C})$ be the C^* -algebra of all $n \times n$ matrices over the complex field.

In their 1992 LAA paper *Generalized Choi Maps in Three Dimensional Matrix Algebra*, Cho, Kye and Lee constructed a class of positive linear maps in $B(M_3(\mathbb{C}), M_3(\mathbb{C}))$:

A Corollary for Generalized Choi Maps in Three Dimensional Matrix Algebra

Let $M_n(\mathbb{C})$ be the C^* -algebra of all $n \times n$ matrices over the complex field.

In their 1992 LAA paper *Generalized Choi Maps in Three Dimensional Matrix Algebra*, Cho, Kye and Lee constructed a class of positive linear maps in $B(M_3(\mathbb{C}), M_3(\mathbb{C}))$:

For nonnegative real numbers a, b and c , the generalized Choi map $\Phi[a, b, c]$ is defined by

$$\Phi[a, b, c](X) = \begin{pmatrix} ax_{11} + bx_{22} + cx_{33} & -x_{12} & -x_{13} \\ -x_{21} & cx_{11} + ax_{22} + bx_{33} & -x_{23} \\ -x_{31} & -x_{32} & bx_{11} + cx_{22} + ax_{33} \end{pmatrix}$$

for $X = [x_{ij}] \in M_3(\mathbb{C})$.

A Corollary for Generalized Choi Maps in Three Dimensional Matrix Algebra

In that paper, conditions on a, b, c were determined for the generalized Choi map $\Phi[a, b, c]$ to be positive, 2-positive, 2-copositive, completely positive, completely copositive and decomposable, respectively.

A Corollary for Generalized Choi Maps in Three Dimensional Matrix Algebra

In that paper, conditions on a, b, c were determined for the generalized Choi map $\Phi[a, b, c]$ to be positive, 2-positive, 2-copositive, completely positive, completely copositive and decomposable, respectively.

Moreover, it was shown that

A Corollary for Generalized Choi Maps in Three Dimensional Matrix Algebra

In that paper, conditions on a, b, c were determined for the generalized Choi map $\Phi[a, b, c]$ to be positive, 2-positive, 2-copositive, completely positive, completely copositive and decomposable, respectively.

Moreover, it was shown that

A Corollary

If the linear map $\Phi[a, b, c]$ is 2-positive or 2-copositive, then it is decomposable.

A Conjecture

A Conjecture

Conjecture 1

Every 2-positive (respectively 2-copositive) map in $B(M_3(\mathbb{C}), M_3(\mathbb{C}))$ is decomposable.

Strong Evidence that All PPTES in $\mathbb{C}^3 \otimes \mathbb{C}^3$ (Two Qutrits) have Schmidt Number 2.

Let ρ be the density matrix for a quantum state in a bipartite system $\mathcal{H}_A \otimes \mathcal{H}_B$. The *Schmidt number* of the density matrix (or the state) ρ is defined by

$$SN(\rho) = \min \left\{ \max_k SR(z_k) \right\},$$

where the minimum is taken over all possible decompositions

$$\rho = \sum_k p_k \cdot z_k z_k^*$$

with z_k being vectors in $\mathcal{H}_A \otimes \mathcal{H}_B$ and $p_k > 0$, $\sum_k p_k = 1$.

Strong Evidence that All PPTES in $\mathbb{C}^3 \otimes \mathbb{C}^3$ (Two Qutrits) have Schmidt Number 2.

In the 2001 *Phy Rev A* paper *Schmidt number witnesses and bound entanglement*, Sanpera, Bruß and Lewenstein showed:

Strong Evidence that All PPTES in $\mathbb{C}^3 \otimes \mathbb{C}^3$ (Two Qutrits) have Schmidt Number 2.

In the 2001 *Phy Rev A* paper *Schmidt number witnesses and bound entanglement*, Sanpera, Bruß and Lewenstein showed:

Validity for some special cases

All positive partial transpose (PPT) entangled states of rank 4 have Schmidt number 2.

Strong Evidence that All PPTES in $\mathbb{C}^3 \otimes \mathbb{C}^3$ (Two Qutrits) have Schmidt Number 2.

In the 2001 *Phy Rev A* paper *Schmidt number witnesses and bound entanglement*, Sanpera, Bruß and Lewenstein showed:

Validity for some special cases

All positive partial transpose (PPT) entangled states of rank 4 have Schmidt number 2.

They also formulated the following conjecture and presented strong evidence of its validity heuristically.

Strong Evidence that All PPTES in $\mathbb{C}^3 \otimes \mathbb{C}^3$ (Two Qutrits) have Schmidt Number 2.

In the 2001 Phy Rev A paper *Schmidt number witnesses and bound entanglement*, Sanpera, Bruß and Lewenstein showed:

Validity for some special cases

All positive partial transpose (PPT) entangled states of rank 4 have Schmidt number 2.

They also formulated the following conjecture and presented strong evidence of its validity heuristically.

Conjecture 2

In $\mathbb{C}^3 \otimes \mathbb{C}^3$, all PPT entangled states have Schmidt number 2.

Dual Cone Relations

There is a classical result on the dual cone relations between quantum states and positive maps pointed out and developed by Størmer, Itoh, Eom and Kye in a series of papers.

Dual Cone Relations

There is a classical result on the dual cone relations between quantum states and positive maps pointed out and developed by Størmer, Itoh, Eom and Kye in a series of papers.

Let us consider the duality between the space $M_m(\mathbb{C}) \otimes M_n(\mathbb{C})$ and the space $B(M_m(\mathbb{C}), M_n(\mathbb{C}))$. Let E_{ij} be the canonical matrix units in $M_m(\mathbb{C})$. For $A = \sum_{i,j=1}^m E_{ij} \otimes A_{ij} \in M_m(\mathbb{C}) \otimes M_n(\mathbb{C})$ and a linear map $\phi \in B(M_m(\mathbb{C}), M_n(\mathbb{C}))$, define a bilinear form:

$$\langle A, \phi \rangle = \sum_{i,j=1}^m \text{Tr}(\phi(E_{ij})A_{ij}^t) = \text{Tr}(A[\phi(E_{ij})]^t).$$

Dual Cone Relations

Denote by $\mathbb{P}_k[m, n]$ and $\mathbb{P}^k[m, n]$ the set of all k -positive maps and the set of all k -copositive maps in $B(M_m(\mathbb{C}), M_n(\mathbb{C}))$, respectively.

Dual Cone Relations

Denote by $\mathbb{P}_k[m, n]$ and $\mathbb{P}^k[m, n]$ the set of all k -positive maps and the set of all k -copositive maps in $B(M_m(\mathbb{C}), M_n(\mathbb{C}))$, respectively.

Define convex cones $\mathbb{V}_k[m, n]$ and $\mathbb{V}^k[m, n]$ in $M_m(\mathbb{C}) \otimes M_n(\mathbb{C})$ as

$$\mathbb{V}_k[m, n] = \{zz^* : SR(z) \leq k, z \text{ in } \mathbb{C}^m \otimes \mathbb{C}^n\}^{\circ\circ},$$

$$\mathbb{V}^k[m, n] = \{(zz^*)^\tau : SR(z) \leq k, z \text{ in } \mathbb{C}^m \otimes \mathbb{C}^n\}^{\circ\circ}.$$

Here τ is partial transposition that acts as transposition only on the first part of a tensor product.

Dual Cone Relations

With the aforementioned notations, by the dual correspondence between maps and states, the following diagram holds:

Dual Cone Relations

With the aforementioned notations, by the dual correspondence between maps and states, the following diagram holds:

Dual Cone Relations

$$\begin{array}{ccccccc}
 \mathbb{V}_1 & \subsetneq & \cdots & \mathbb{V}_k & \subsetneq & \mathbb{V}_{m \wedge n} & = (M_m(\mathbb{C}) \otimes M_n(\mathbb{C}))^+ \\
 \updownarrow & & & \updownarrow & & \updownarrow & \\
 \mathbb{P}_1 & \supsetneq & \cdots & \mathbb{P}_k & \supsetneq & \mathbb{P}_{m \wedge n} & \cong (M_m(\mathbb{C}) \otimes M_n(\mathbb{C}))^+
 \end{array}$$

Dual Cone Relations

With the aforementioned notations, by the dual correspondence between maps and states, the following diagram holds:

Dual Cone Relations

$$\begin{array}{ccccccc}
 \mathbb{V}_1 & \subsetneq & \cdots & \mathbb{V}_k & \subsetneq & \mathbb{V}_{m \wedge n} & = (M_m(\mathbb{C}) \otimes M_n(\mathbb{C}))^+ \\
 \updownarrow & & & \updownarrow & & \updownarrow & \\
 \mathbb{P}_1 & \supsetneq & \cdots & \mathbb{P}_k & \supsetneq & \mathbb{P}_{m \wedge n} & \cong (M_m(\mathbb{C}) \otimes M_n(\mathbb{C}))^+
 \end{array}$$

where $m \wedge n = \min\{m, n\}$, and a similar diagram holds in case of copositivity.

Dual Cone Relations when $m = n = 3$

Denote by \mathbb{D} the cone of all decomposable maps and \mathbb{T} the cone of all positive partial transpose states.

Dual Cone Relations when $m = n = 3$

Denote by \mathbb{D} the cone of all decomposable maps and \mathbb{T} the cone of all positive partial transpose states.

In Kye's paper *Facial structures for various notions of positivity and applications to the theory of entanglement*, the two conjectures are unified in the dual cone scheme.

Dual Cone Relations when $m = n = 3$

Denote by \mathbb{D} the cone of all decomposable maps and \mathbb{T} the cone of all positive partial transpose states.

In Kye's paper *Facial structures for various notions of positivity and applications to the theory of entanglement*, the two conjectures are unified in the dual cone scheme.

Dual Cone Relations when $m = n = 3$

$$\begin{array}{ccccccc}
 \text{Conj2 : } & \mathbb{V}_1 & \subsetneq & \mathbb{T}(?) & \subsetneq & \mathbb{V}_2 & \subsetneq & \mathbb{V}_3 = (M_3(\mathbb{C}) \otimes M_3(\mathbb{C}))^+ \\
 & \updownarrow & & \updownarrow & & \updownarrow & & \updownarrow \\
 \text{Conj1 : } & \mathbb{P}_1 & \supsetneq & \mathbb{D}(?) & \supsetneq & \mathbb{P}_2 & \supsetneq & \mathbb{P}_3 \cong (M_3(\mathbb{C}) \otimes M_3(\mathbb{C}))^+
 \end{array}$$

A Peel-off Theorem

An astonishing result first appeared in Marciniak's paper *On extremal positive maps acting between type I factors*.

A Peel-off Theorem

An astonishing result first appeared in Marciniak's paper *On extremal positive maps acting between type I factors*.

Peel-off Theorem (Marciniak)

If ϕ is a non-zero 2-positive map, then there exists a non-zero completely positive map ψ such that $\phi \geq \psi$.

Trivial Lifting

We will present a slightly stronger version (Choi Decomposition) of the peel-off result by block-matrix approach, which was shown by Choi for the case of 2-positive maps.

Trivial Lifting

We will present a slightly stronger version (Choi Decomposition) of the peel-off result by block-matrix approach, which was shown by Choi for the case of 2-positive maps.

Definition of Trivial Lifting

Given a linear map $\chi \in B(M_s(\mathbb{C}), M_n(\mathbb{C}))$, fix the canonical matrix unit basis E_{ij} , $i, j = 1, \dots, s$, in $M_s(\mathbb{C})$, under which the Choi matrix is $C_\chi = [\chi(E_{ij})]_{i,j=1}^s \in M_s(M_n(\mathbb{C}))$. Given

$L = \{n_1, \dots, n_p\} \subset \{1, \dots, s+p\}$, where $n_1 < \dots < n_p$, extend the matrix C_χ to a $(s+p) \times (s+p)$ block matrix

$C_L^{lift} \in M_{s+p}(M_n(\mathbb{C}))$ by adding one row and one column of $n \times n$ zero matrices at the n_k^{th} level for each $k = 1, \dots, p$ as follows:

Trivial Lifting

Definition of Trivial Lifting

$$C_L^{lift} \triangleq \begin{matrix} & \begin{matrix} 1^{st} & \cdots & n_k^{th} & \cdots & (s+p)^{th} \end{matrix} \\ \begin{matrix} 1^{st} \\ \vdots \\ n_k^{th} \\ \vdots \\ (s+p)^{th} \end{matrix} & \begin{pmatrix} \chi(E_{11}) & \cdots & 0 & \cdots & \chi(E_{1,s}) \\ \vdots & \ddots & 0 & \ddots & \vdots \\ 0 & 0 & 0 & 0 & 0 \\ \vdots & \ddots & 0 & \ddots & \vdots \\ \chi(E_{s,1}) & \cdots & 0 & \cdots & \chi(E_{s,s}) \end{pmatrix} \end{matrix}.$$

Denote by $\tilde{\chi}_L$ the map in $B(M_{s+p}(\mathbb{C}), M_n(\mathbb{C}))$ associated with the Choi matrix $C_{\tilde{\chi}_L} = [\tilde{\chi}_L(E_{ij})]_{i,j=1}^{s+p} = C_L^{lift}$. Then the map $\tilde{\chi}_L$ is called a L -trivial lifting of the original map χ . If $L = \{q\}$ is a singleton, simply denote by $\tilde{\chi}_q$ the q -trivial lifting of χ .

Trivial Lifting

Remarks for the trivial lifting:

Trivial Lifting

Remarks for the trivial lifting:

Remark 1 for trivial lifting

A map χ is k -positive (respectively k -copositive) if and only if its trivial lifting $\tilde{\chi}_L$ is k -positive (respectively k -copositive).

Trivial Lifting

Remarks for the trivial lifting:

Remark 1 for trivial lifting

A map χ is k -positive (respectively k -copositive) if and only if its trivial lifting $\tilde{\chi}_L$ is k -positive (respectively k -copositive).

Remark 2 for trivial lifting

A map χ is decomposable if and only if its trivial lifting $\tilde{\chi}_L$ is decomposable.

Main Result

Motivated by Choi's block matrix approach regarding the peel-off theorem, we obtain the following result:

Main Result

Motivated by Choi's block matrix approach regarding the peel-off theorem, we obtain the following result:

Theorem 1 (Choi Decomposition Theorem)

Let ϕ be a non-zero k -positive ($2 \leq k < \min\{m, n\}$) map in $B(M_m(\mathbb{C}), M_n(\mathbb{C}))$. Then there exists a decomposition $\phi = \psi + \gamma$, where ψ is a non-zero completely positive map and γ is a p -trivial lifting of a $(k-1)$ -positive map in $B(M_{m-1}(\mathbb{C}), M_n(\mathbb{C}))$, for some $p \in \{1, \dots, m\}$.

Main Result

Motivated by Choi's block matrix approach regarding the peel-off theorem, we obtain the following result:

Theorem 1 (Choi Decomposition Theorem)

Let ϕ be a non-zero k -positive ($2 \leq k < \min\{m, n\}$) map in $B(M_m(\mathbb{C}), M_n(\mathbb{C}))$. Then there exists a decomposition $\phi = \psi + \gamma$, where ψ is a non-zero completely positive map and γ is a p -trivial lifting of a $(k-1)$ -positive map in $B(M_{m-1}(\mathbb{C}), M_n(\mathbb{C}))$, for some $p \in \{1, \dots, m\}$.

Notice that the dimension of the space where the remaining map γ resides is reduced.

Sketch of the Proof: Useful Lemmas

Lemma 1: Positivity in terms of Block Matrix

Suppose a hermitian matrix M is partitioned as

$$M = \begin{pmatrix} A & B \\ B^* & C \end{pmatrix},$$

where A and C are square matrices. TFAE:

- 1 $M \geq 0$,
- 2 $A \geq 0$, $M/A = C - B^*A^\dagger B \geq 0$, $\text{range}(B) \subset \text{range}(A)$,
- 3 $C \geq 0$, $M/C = A - BC^\dagger B^* \geq 0$, $\text{range}(B^*) \subset \text{range}(C)$.

Here A^\dagger and C^\dagger refer to the Moore-Penrose pseudo inverses of A and C , respectively.

Sketch of the Proof: Useful Lemmas

Lemma 2: Properties of the Moore-Penrose Pseudo Inverse

- ① $AA^\dagger A = A$, $A^\dagger AA^\dagger = A^\dagger$.
- ② $(AA^\dagger)^* = AA^\dagger$, $(A^\dagger A)^* = A^\dagger A$.
- ③ AA^\dagger is the orthogonal projector onto the range of A , $A^\dagger A$ is the orthogonal projector onto the range of A^* .
- ④ If A is invertible, then $A^\dagger = A^{-1}$.
- ⑤ If $A \geq 0$, then $A^\dagger \geq 0$.

Sketch of the Proof: Block Matrix Approach

Let us look at the Choi matrix C_ϕ for ϕ , with
 $A_{ij} = \phi(E_{ij})$, $i, j = 1, \dots, m$.

Sketch of the Proof: Block Matrix Approach

Let us look at the Choi matrix C_ϕ for ϕ , with
 $A_{ij} = \phi(E_{ij})$, $i, j = 1, \dots, m$.

Choi Decomposition: Original Part

$$C_\phi = \begin{pmatrix} A_{11} & \cdots & A_{1j} & \cdots & A_{1m} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ A_{i1} & \cdots & A_{ij} & \cdots & A_{im} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ A_{m1} & \cdots & A_{mj} & \cdots & A_{mm} \end{pmatrix}$$

Sketch of the Proof: Block Matrix Approach

Let us look at the Choi matrix C_ϕ for ϕ , with
 $A_{ij} = \phi(E_{ij})$, $i, j = 1, \dots, m$.

Choi Decomposition: Original Part

$$C_\phi = \begin{pmatrix} A_{11} & \cdots & A_{1j} & \cdots & A_{1m} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ A_{i1} & \cdots & A_{ij} & \cdots & A_{im} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ A_{m1} & \cdots & A_{mj} & \cdots & A_{mm} \end{pmatrix}$$

Observation 1: WOLOG, assume that $\phi(E_{mm}) \neq 0$.

Sketch of the Proof: Block Matrix Approach

The peel-off part is a matrix with $A_{im}A_{mm}^\dagger A_{mj}$ in its (i, j) -entry.

Sketch of the Proof: Block Matrix Approach

The peel-off part is a matrix with $A_{im}A_{mm}^\dagger A_{mj}$ in its (i, j) -entry.

Choi Decomposition: Peel-off Part

$$U = \begin{pmatrix} A_{1m}A_{mm}^\dagger A_{m1} & \cdots & A_{1m}A_{mm}^\dagger A_{mj} & \cdots & A_{1m}A_{mm}^\dagger A_{mm} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ A_{im}A_{mm}^\dagger A_{m1} & \cdots & A_{im}A_{mm}^\dagger A_{mj} & \cdots & A_{im}A_{mm}^\dagger A_{mm} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ A_{mm}A_{mm}^\dagger A_{m1} & \cdots & A_{mm}A_{mm}^\dagger A_{mj} & \cdots & A_{mm}A_{mm}^\dagger A_{mm} \end{pmatrix}$$

Sketch of the Proof: Block Matrix Approach

The peel-off part is a matrix with $A_{im}A_{mm}^\dagger A_{mj}$ in its (i, j) -entry.

Choi Decomposition: Peel-off Part

$$U = \begin{pmatrix} A_{1m}A_{mm}^\dagger A_{m1} & \cdots & A_{1m}A_{mm}^\dagger A_{mj} & \cdots & A_{1m}A_{mm}^\dagger A_{mm} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ A_{im}A_{mm}^\dagger A_{m1} & \cdots & A_{im}A_{mm}^\dagger A_{mj} & \cdots & A_{im}A_{mm}^\dagger A_{mm} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ A_{mm}A_{mm}^\dagger A_{m1} & \cdots & A_{mm}A_{mm}^\dagger A_{mj} & \cdots & A_{mm}A_{mm}^\dagger A_{mm} \end{pmatrix}$$

Observation 2: $U \geq 0$, and U is non-zero.

Sketch of the Proof: Block Matrix Approach

The remaining part is a matrix with $R_{ij} = A_{ij} - A_{im}A_{mm}^\dagger A_{mj}$ in its (i, j) -entry.

Sketch of the Proof: Block Matrix Approach

The remaining part is a matrix with $R_{ij} = A_{ij} - A_{im}A_{mm}^\dagger A_{mj}$ in its (i, j) -entry.

Choi Decomposition: Remaining Part

$$R = \begin{pmatrix} A_{11} - A_{1m}A_{mm}^\dagger A_{m1} & \cdots & A_{1j} - A_{1m}A_{mm}^\dagger A_{mj} & \cdots & A_{1m} - A_{1m}A_{mm}^\dagger A_{mm} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ A_{i1} - A_{im}A_{mm}^\dagger A_{m1} & \cdots & A_{ij} - A_{im}A_{mm}^\dagger A_{mj} & \cdots & A_{im} - A_{im}A_{mm}^\dagger A_{mm} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ A_{m1} - A_{mm}A_{mm}^\dagger A_{m1} & \ddots & A_{mj} - A_{mm}A_{mm}^\dagger A_{mj} & \cdots & A_{mm} - A_{mm}A_{mm}^\dagger A_{mm} \end{pmatrix}$$

Sketch of the Proof: Block Matrix Approach

The remaining part is a matrix with $R_{ij} = A_{ij} - A_{im}A_{mm}^\dagger A_{mj}$ in its (i, j) -entry.

Choi Decomposition: Remaining Part

$$R = \begin{pmatrix} A_{11} - A_{1m}A_{mm}^\dagger A_{m1} & \cdots & A_{1j} - A_{1m}A_{mm}^\dagger A_{mj} & \cdots & A_{1m} - A_{1m}A_{mm}^\dagger A_{mm} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ A_{i1} - A_{im}A_{mm}^\dagger A_{m1} & \cdots & A_{ij} - A_{im}A_{mm}^\dagger A_{mj} & \cdots & A_{im} - A_{im}A_{mm}^\dagger A_{mm} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ A_{m1} - A_{mm}A_{mm}^\dagger A_{m1} & \ddots & A_{mj} - A_{mm}A_{mm}^\dagger A_{mj} & \cdots & A_{mm} - A_{mm}A_{mm}^\dagger A_{mm} \end{pmatrix}$$

Observation 3: Entries in last row and last column of R are zero matrices

Sketch of the Proof: Block Matrix Approach

Now $C_\phi = U + R = C_\psi + C_\gamma$, with ψ completely positive.

Sketch of the Proof: Block Matrix Approach

Now $C_\phi = U + R = C_\psi + C_\gamma$, with ψ completely positive.

Question: What will γ be?

Sketch of the Proof: Block Matrix Approach

Now $C_\phi = U + R = C_\psi + C_\gamma$, with ψ completely positive.

Question: What will γ be?

Good News: k -positivity of ϕ guarantees $(k - 1)$ -positivity of γ .

Sketch of the Proof: Block Matrix Approach

Now $C_\phi = U + R = C_\psi + C_\gamma$, with ψ completely positive.

Question: What will γ be?

Good News: k -positivity of ϕ guarantees $(k - 1)$ -positivity of γ .

Choi Decomposition: Employ k -positivity of ϕ for $\xi\xi^*$

$$\xi\xi^* = \begin{pmatrix} w^1(w^1)^* & \cdots & w^1(w^j)^* & \cdots & w^1 e_m^* \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ w^i(w^1)^* & \cdots & w^i(w^j)^* & \cdots & w^i e_m^* \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ e_m(w^1)^* & \cdots & e_m(w^j)^* & \cdots & e_m e_m^* \end{pmatrix} \geq 0$$

Here $\xi = [w^1; \dots; w^{k-1}; e_m]$, where $w^1, w^2, \dots, w^{k-1} \in \mathbb{C}^m$ are arbitrary column vectors, and $e_m = (0, \dots, 0, 1)^T \in \mathbb{C}^m$.

Sketch of the Proof: Block Matrix Approach

Choi Decomposition: Employ k -positivity of ϕ for $\xi\xi^*$

$$(id_k \otimes \phi)(\xi\xi^*) = \begin{pmatrix} \phi(w^1(w^1)^*) & \cdots & \phi(w^1(w^j)^*) & \cdots & \phi(w^1 e_m^*) \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ \phi(w^i(w^1)^*) & \cdots & \phi(w^i(w^j)^*) & \cdots & \phi(w^i e_m^*) \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ \phi(e_m(w^1)^*) & \cdots & \phi(e_m(w^j)^*) & \cdots & \phi(e_m e_m^*) \end{pmatrix} \geq 0.$$

Sketch of the Proof: Block Matrix Approach

Choi Decomposition: Employ k -positivity of ϕ for $\xi\xi^*$

$$(id_k \otimes \phi)(\xi\xi^*) = \begin{pmatrix} \phi(w^1(w^1)^*) & \cdots & \phi(w^1(w^j)^*) & \cdots & \phi(w^1 e_m^*) \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ \phi(w^i(w^1)^*) & \cdots & \phi(w^i(w^j)^*) & \cdots & \phi(w^i e_m^*) \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ \phi(e_m(w^1)^*) & \cdots & \phi(e_m(w^j)^*) & \cdots & \phi(e_m e_m^*) \end{pmatrix} \geq 0.$$

Observation 4: Recall Lemma 1.

Sketch of the Proof: Block Matrix Approach

By equivalence of Condition 1 and Condition 3 in Lemma 1, the condition $(id_k \otimes \phi)(\xi\xi^*) \geq 0$ expands to:

Sketch of the Proof: Block Matrix Approach

By equivalence of Condition 1 and Condition 3 in Lemma 1, the condition $(id_k \otimes \phi)(\xi\xi^*) \geq 0$ expands to:

Choi Decomposition: Employ k -positivity of ϕ for $\xi\xi^*$

$$\begin{pmatrix} \phi(w^1(w^1)^*) & \cdots & \phi(w^1(w^{k-1})^*) \\ \vdots & \ddots & \vdots \\ \phi(w^{k-1}(w^1)^*) & \cdots & \phi(w^{k-1}(w^{k-1})^*) \end{pmatrix} \geq \begin{pmatrix} \phi(w^1 e_m^*) \\ \vdots \\ \phi(w^{k-1} e_m^*) \end{pmatrix} \phi(e_m e_m^*)^\dagger \left(\phi(e_m(w^1)^*) \quad \cdots \quad \phi(e_m(w^{k-1})^*) \right).$$

Sketch of the Proof: Block Matrix Approach

By equivalence of Condition 1 and Condition 3 in Lemma 1, the condition $(id_k \otimes \phi)(\xi\xi^*) \geq 0$ expands to:

Choi Decomposition: Employ k -positivity of ϕ for $\xi\xi^*$

$$\begin{pmatrix} \phi(w^1(w^1)^*) & \cdots & \phi(w^1(w^{k-1})^*) \\ \vdots & \ddots & \vdots \\ \phi(w^{k-1}(w^1)^*) & \cdots & \phi(w^{k-1}(w^{k-1})^*) \end{pmatrix} \geq \begin{pmatrix} \phi(w^1 e_m^*) \\ \vdots \\ \phi(w^{k-1} e_m^*) \end{pmatrix} \phi(e_m e_m^*)^\dagger \left(\phi(e_m(w^1)^*) \quad \cdots \quad \phi(e_m(w^{k-1})^*) \right).$$

Observation 5: The (s, t) -entry of above RHS is $\psi(w^s(w^t)^*)$:

Sketch of the Proof: Block Matrix Approach

Choi Decomposition: Employ k -positivity of ϕ for $\xi\xi^*$

$$\begin{aligned}
 & \phi(w^s e_m^*) \phi(e_m e_m^*)^\dagger \phi(e_m (w^t)^*) \\
 &= \left(\sum_{i=1}^m w_i^s \phi(E_{im}) \right) \phi(E_{mm})^\dagger \left(\sum_{j=1}^m \overline{w_j^t} \phi(E_{mj}) \right) \\
 &= \sum_{i=1}^m \sum_{j=1}^m w_i^s \overline{w_j^t} \left(\phi(E_{im}) \phi(E_{mm})^\dagger \phi(E_{mj}) \right) \\
 &= \sum_{i=1}^m \sum_{j=1}^m w_i^s \overline{w_j^t} (A_{im} A_{mm}^\dagger A_{mj}) \\
 &= \sum_{i=1}^m \sum_{j=1}^m w_i^s \overline{w_j^t} \psi(e_i e_j^*) \\
 &= \psi(w^s (w^t)^*)
 \end{aligned}$$

Sketch of the Proof: Block Matrix Approach

This proves that $\gamma = \phi - \psi$ is $(k - 1)$ -positive.

Sketch of the Proof: Block Matrix Approach

This proves that $\gamma = \phi - \psi$ is $(k - 1)$ -positive.

Choi Decomposition: Employ k -positivity of ϕ for $\xi\xi^*$

$$\begin{pmatrix} \gamma(w^1(w^1)^*) & \dots & \gamma(w^1(w^{k-1})^*) \\ \vdots & \ddots & \vdots \\ \gamma(w^{k-1}(w^1)^*) & \dots & \gamma(w^{k-1}(w^{k-1})^*) \end{pmatrix} \geq 0, \forall w^1, \dots, w^{k-1} \in \mathbb{C}^m.$$

Sketch of the Proof: Block Matrix Approach

This proves that $\gamma = \phi - \psi$ is $(k - 1)$ -positive.

Choi Decomposition: Employ k -positivity of ϕ for $\xi\xi^*$

$$\begin{pmatrix} \gamma(w^1(w^1)^*) & \dots & \gamma(w^1(w^{k-1})^*) \\ \vdots & \ddots & \vdots \\ \gamma(w^{k-1}(w^1)^*) & \dots & \gamma(w^{k-1}(w^{k-1})^*) \end{pmatrix} \geq 0, \forall w^1, \dots, w^{k-1} \in \mathbb{C}^m.$$

Combining Observation 3 and the above fact, we know the form of the remaining map γ .

Sketch of the Proof: Block Matrix Approach

Denote the matrix $R = C_\gamma$ by:

$$R = \begin{pmatrix} \mathbf{K} & 0 \\ \vdots & \vdots \\ 0 & \dots & 0 \end{pmatrix} = \begin{pmatrix} \mathbf{C}_K & 0 \\ \vdots & \vdots \\ 0 & \dots & 0 \end{pmatrix}.$$

Sketch of the Proof: Block Matrix Approach

Denote the matrix $R = C_\gamma$ by:

$$R = \begin{pmatrix} \mathbf{K} & 0 \\ & \vdots \\ 0 & \dots & 0 \end{pmatrix} = \begin{pmatrix} \mathbf{C}_\kappa & 0 \\ & \vdots \\ 0 & \dots & 0 \end{pmatrix}.$$

Choi Decomposition: γ is a trivial-lifting of κ

The map $\kappa \in B(M_{m-1}(\mathbb{C}), M_n(\mathbb{C}))$ is defined by its Choi matrix $C_\kappa = K \in M_{m-1}(M_n(\mathbb{C}))$ through $\kappa(E_{st}) = K_{st}$, $s, t = 1, \dots, m-1$. It is obvious that $\gamma \in B(M_m(\mathbb{C}), M_n(\mathbb{C}))$ is the m -trivial lifting of $\kappa \in B(M_{m-1}(\mathbb{C}), M_n(\mathbb{C}))$.

Sketch of the Proof: Block Matrix Approach

Denote the matrix $R = C_\gamma$ by:

$$R = \begin{pmatrix} \mathbf{K} & 0 \\ \vdots & \vdots \\ 0 & \dots & 0 \end{pmatrix} = \begin{pmatrix} \mathbf{C}_\kappa & 0 \\ \vdots & \vdots \\ 0 & \dots & 0 \end{pmatrix}.$$

Choi Decomposition: γ is a trivial-lifting of κ

The map $\kappa \in B(M_{m-1}(\mathbb{C}), M_n(\mathbb{C}))$ is defined by its Choi matrix $C_\kappa = K \in M_{m-1}(M_n(\mathbb{C}))$ through $\kappa(E_{st}) = K_{st}$, $s, t = 1, \dots, m-1$. It is obvious that $\gamma \in B(M_m(\mathbb{C}), M_n(\mathbb{C}))$ is the m -trivial lifting of $\kappa \in B(M_{m-1}(\mathbb{C}), M_n(\mathbb{C}))$.

A similar result holds for k -copositive maps.

Choi Decomposition

Applying Theorem 1 repeatedly,

Theorem 2

Let $2 \leq k < \min\{m, n\}$. Any non-zero k -positive (respectively k -copositive) map in $B(M_m(\mathbb{C}), M_n(\mathbb{C}))$ is the sum of at most $(k - 1)$ many non-zero completely positive (respectively completely copositive) maps and a positive map which is the trivial lifting of a positive map in $B(M_{m-k+1}(\mathbb{C}), M_n(\mathbb{C}))$.

Choi Decomposition

Applying Theorem 1 repeatedly,

Theorem 2

Let $2 \leq k < \min\{m, n\}$. Any non-zero k -positive (respectively k -copositive) map in $B(M_m(\mathbb{C}), M_n(\mathbb{C}))$ is the sum of at most $(k - 1)$ many non-zero completely positive (respectively completely copositive) maps and a positive map which is the trivial lifting of a positive map in $B(M_{m-k+1}(\mathbb{C}), M_n(\mathbb{C}))$.

Remark: The Choi decomposition may no longer be valid for a general positive map ϕ , even when ϕ is in $B(M_2(\mathbb{C}), M_2(\mathbb{C}))$.

An Affirmative Answer to the Conjecture

Theorem 3

Every 2-positive or 2-copositive map ϕ in $B(M_3(\mathbb{C}), M_3(\mathbb{C}))$ is decomposable.

An Affirmative Answer to the Conjecture

Theorem 3

Every 2-positive or 2-copositive map ϕ in $B(M_3(\mathbb{C}), M_3(\mathbb{C}))$ is decomposable.

Proof: WOLOG, we assume the 2-positive (respectively 2-copositive) map ϕ is not zero. In this concrete case of $B(M_3(\mathbb{C}), M_3(\mathbb{C}))$, the peel-off process yields that:

$$\phi = \psi + \tilde{\kappa}_p \text{ for some } p \in \{1, \dots, m\},$$

where ψ is completely positive (respectively completely copositive) and $\tilde{\kappa}_p$ is a p -trivial lifting of a positive map $\kappa \in B(M_2(\mathbb{C}), M_3(\mathbb{C}))$.

An Affirmative Answer to the Conjecture

We will use an important result of Størmer and Woronowicz:

An Affirmative Answer to the Conjecture

We will use an important result of Størmer and Woronowicz:

Positive=Decomposable in $2 \otimes 2$, $2 \otimes 3$ and $3 \otimes 2$.

$\mathbb{P}_1[m, n] = \mathbb{D}[m, n]$ holds when $mn \leq 6$.

An Affirmative Answer to the Conjecture

We will use an important result of Størmer and Woronowicz:

Positive=Decomposable in $2 \otimes 2$, $2 \otimes 3$ and $3 \otimes 2$.

$\mathbb{P}_1[m, n] = \mathbb{D}[m, n]$ holds when $mn \leq 6$.

Since every positive map in $B(M_2(\mathbb{C}), M_3(\mathbb{C}))$ is decomposable in $B(M_2(\mathbb{C}), M_3(\mathbb{C}))$, by properties of trivial lifting, the lifted map $\tilde{\kappa}_p$ is decomposable in $B(M_3(\mathbb{C}), M_3(\mathbb{C}))$.

An Affirmative Answer to the Conjecture

We will use an important result of Størmer and Woronowicz:

Positive=Decomposable in $2 \otimes 2$, $2 \otimes 3$ and $3 \otimes 2$.

$\mathbb{P}_1[m, n] = \mathbb{D}[m, n]$ holds when $mn \leq 6$.

Since every positive map in $B(M_2(\mathbb{C}), M_3(\mathbb{C}))$ is decomposable in $B(M_2(\mathbb{C}), M_3(\mathbb{C}))$, by properties of trivial lifting, the lifted map $\tilde{\kappa}_p$ is decomposable in $B(M_3(\mathbb{C}), M_3(\mathbb{C}))$.

Hence, $\phi = \psi + \tilde{\kappa}_p$ is also decomposable.

A Corollary & An Example

Corollary 4

Every indecomposable map in $B(M_3(\mathbb{C}), M_3(\mathbb{C}))$ is atomic (i.e., *not* the sum of a 2-positive map and a 2-copositive map).

A Corollary & An Example

Corollary 4

Every indecomposable map in $B(M_3(\mathbb{C}), M_3(\mathbb{C}))$ is atomic (i.e., *not* the sum of a 2-positive map and a 2-copositive map).

Remark: There exist different methods to decompose the 2-positive generalized Choi map $\Phi[a, b, c]$ into a sum of a completely positive map and a completely copositive map.

A Corollary & An Example

Corollary 4

Every indecomposable map in $B(M_3(\mathbb{C}), M_3(\mathbb{C}))$ is atomic (i.e., *not* the sum of a 2-positive map and a 2-copositive map).

Remark: There exist different methods to decompose the 2-positive generalized Choi map $\Phi[a, b, c]$ into a sum of a completely positive map and a completely copositive map.

An Example

$$\Phi[a, b, c](X) = \begin{pmatrix} ax_{11} + bx_{22} + cx_{33} & -x_{12} & -x_{13} \\ -x_{21} & cx_{11} + ax_{22} + bx_{33} & -x_{23} \\ -x_{31} & -x_{32} & bx_{11} + cx_{22} + ax_{33} \end{pmatrix}$$

for $X = [x_{ij}] \in M_3(\mathbb{C})$. Here $a \in [1, 2)$ and $bc \geq (2 - a)(b + c)$.

A Corollary & An Example

An Example: Decomposition 1

$\Phi[a, b, c] = \Phi_1 + \Phi_2$, where

$$\Phi_1 \begin{bmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{bmatrix} = \begin{bmatrix} ax_{11} + bx_{22} + cx_{33} & -x_{12} & -x_{13} \\ -x_{21} & cx_{11} + ax_{22} & (\frac{2}{a} - a)x_{23} \\ -x_{31} & (\frac{2}{a} - a)x_{32} & bx_{11} + ax_{33} \end{bmatrix}, (CP)$$

$$\Phi_2 \begin{bmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & bx_{33} & (a - 1 - \frac{2}{a})x_{23} \\ 0 & (a - 1 - \frac{2}{a})x_{32} & cx_{22} \end{bmatrix} (CcoP).$$

A Corollary & An Example

An Example: Decomposition 1

$\Phi[a, b, c] = \Phi_1 + \Phi_2$, where

$$\Phi_1 \begin{bmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{bmatrix} = \begin{bmatrix} ax_{11} + bx_{22} + cx_{33} & -x_{12} & -x_{13} \\ -x_{21} & cx_{11} + ax_{22} & (\frac{2}{a} - a)x_{23} \\ -x_{31} & (\frac{2}{a} - a)x_{32} & bx_{11} + ax_{33} \end{bmatrix}, (CP)$$

$$\Phi_2 \begin{bmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & bx_{33} & (a - 1 - \frac{2}{a})x_{23} \\ 0 & (a - 1 - \frac{2}{a})x_{32} & cx_{22} \end{bmatrix} (CcoP).$$

Another decomposition given by Cho, Kye and Lee is:

A Corollary & An Example

An Example: Decomposition 1

$\Phi[a, b, c] = \Phi_1 + \Phi_2$, where

$$\Phi_1 \begin{bmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{bmatrix} = \begin{bmatrix} ax_{11} + bx_{22} + cx_{33} & -x_{12} & -x_{13} \\ -x_{21} & cx_{11} + ax_{22} & (\frac{2}{a} - a)x_{23} \\ -x_{31} & (\frac{2}{a} - a)x_{32} & bx_{11} + ax_{33} \end{bmatrix}, (CP)$$

$$\Phi_2 \begin{bmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & bx_{33} & (a - 1 - \frac{2}{a})x_{23} \\ 0 & (a - 1 - \frac{2}{a})x_{32} & cx_{22} \end{bmatrix} (CcoP).$$

Another decomposition given by Cho, Kye and Lee is:

An Example: Decomposition 2

$$\Phi[a, b, c] = (1 - \sqrt{bc})\Phi\left[\frac{a - \sqrt{bc}}{1 - \sqrt{bc}}, 0, 0\right] (CP) + \sqrt{bc}\Phi\left[1, \sqrt{\frac{b}{c}}, \sqrt{\frac{c}{b}}\right] (CcoP).$$

Question 1: An Algorithm for Decomposition

Given an arbitrary decomposable map, is there a canonical algorithm to decompose it?

Question 1: An Algorithm for Decomposition





Given an arbitrary decomposable map, is there a canonical algorithm to decompose it?

Is such an algorithm possible, even in $B(M_2(\mathbb{C}), M_2(\mathbb{C}))$?





Question 2: An Example in Higher Dimensions

Does there exist a 2-positive but indecomposable map in $B(M_3(\mathbb{C}), M_4(\mathbb{C}))$?





References

-  S. J. Cho, S.-H. Kye and S. G. Lee, *Generalized Choi maps in three-dimensional matrix algebra*, Linear Alg. Appl. **171** (1992), 213-224.
-  M.-D. Choi, Seminar talk at National University of Singapore, 30 July 2015.
-  M.-H. Eom and S.-H. Kye, *Duality for positive linear maps in matrix algebras*, Math. Scan. **86** (2000), 130-142
-  T. Itoh, *Positive maps and cones in C^* -algebras*, Math. Japonica, **31** (1986), 607-616.

References

-  S.-H. Kye, *Facial structures for various notions of positivity and applications to the theory of entanglement*, Rev. Math. Phys. **25** (2013), 1330002.
-  M. Marciniak, *On extremal positive maps acting between type I factors*, Noncommutative Harmonic Analysis with Applications to probability II, Banach Center Publications, **89** (2010), 201-221.
-  H. Osaka, *Indecomposable positive maps in low dimensional matrix algebra*, Linear Alg. Appl. **153** (1991), 73-83.
-  A. Sanpera, D. Bruß and M. Lewenstein, *Schmidt number witnesses and bound entanglement*, Phys Rev. A **63** (2001), 050301.

References

-  E. Størmer, *Positive linear maps of operator algebras*, Acta Mathematica **110** (1963), 233-278.
-  B. M. Terhal and P. Horodecki, *A Schmidt number for density matrices*, Phys Rev. A **61** (2000), 040301.
-  S. L. Woronowicz, *Positive maps of low dimensional matrix algebras*, Rep. Math. Phys. **10** (1976), 165-183.
-  Y. Yang, D. H. Leung and W. S. Tang, *All 2-positive linear maps from $M_3(\mathbb{C})$ to $M_3(\mathbb{C})$ are decomposable*, Linear Alg. Appl. **503** (2016), 233-247.

Thank you for your attention!